

WORLD TRADE ORGANIZATION

G/SPS/N/JPN/302

29 August 2012

(12-4647)

Committee on Sanitary and Phytosanitary Measures

Original: English

NOTIFICATION

1.	Notifying Member: <u>JAPAN</u> If applicable, name of local government involved:
2.	Agency responsible: Ministry of Health, Labour and Welfare
3.	Products covered (provide tariff item number(s) as specified in national schedules deposited with the WTO; ICS numbers should be provided in addition, where applicable): Food additive (Azoxystrobin and Chlorous Acid Water)
4.	Regions or countries likely to be affected, to the extent relevant or practicable: <input checked="" type="checkbox"/> All trading partners <input type="checkbox"/> Specific regions or countries:
5.	Title of the notified document: Amendment to the Enforcement Ordinance of the Food Sanitation Law and the Standards and Specifications for Foods and Food Additives Language(s): English Number of pages: 6 http://members.wto.org/crnattachments/2012/sps/JPN/12_3435_00_e.pdf
6.	Description of content: Authorization of Azoxystrobin and Chlorous Acid Water as food additives and establishment of standards and specifications for this substance.
7.	Objective and rationale: <input checked="" type="checkbox"/> food safety, <input type="checkbox"/> animal health, <input type="checkbox"/> plant protection, <input type="checkbox"/> protect humans from animal/plant pest or disease, <input type="checkbox"/> protect territory from other damage from pests.
8.	Is there a relevant international standard? If so, identify the standard: <input checked="" type="checkbox"/> Codex Alimentarius Commission (<i>e.g. title or serial number of Codex standard or related text</i>) <input type="checkbox"/> World Organization for Animal Health (OIE) (<i>e.g. Terrestrial or Aquatic Animal Health Code, chapter number</i>) <input type="checkbox"/> International Plant Protection Convention (<i>e.g. ISPM number</i>) <input type="checkbox"/> None Does this proposed regulation conform to the relevant international standard? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No If no, describe, whenever possible, how and why it deviates from the international standard:

9.	Other relevant documents and language(s) in which these are available: Food Sanitation Law (available in English). When adopted, this amendment is to be published in Kampo (Official Japanese Governmental Gazette) (available in Japanese).
10.	Proposed date of adoption (dd/mm/yy): As soon as possible after the final date for the comment period. Proposed date of publication (dd/mm/yy): As soon as possible after the final date for the comment period.
11.	Proposed date of entry into force: <input type="checkbox"/> Six months from date of publication, and/or (dd/mm/yy): As soon as possible after publication. <input checked="" type="checkbox"/> Trade facilitating measure
12.	Final date for comments: <input checked="" type="checkbox"/> Sixty days from the date of circulation of the notification and/or (dd/mm/yy): 28 October 2012 Agency or authority designated to handle comments: <input type="checkbox"/> National Notification Authority, <input checked="" type="checkbox"/> National Enquiry Point. Address, fax number and e-mail address (if available) of other body:
13.	Texts available from: <input type="checkbox"/> National Notification Authority, <input checked="" type="checkbox"/> National Enquiry Point. Address, fax number and e-mail address (if available) of other body: Standards Information Service International Trade Division Economic Affairs Bureau Ministry of Foreign Affairs 2-2-1 Kasumigaseki, Chiyoda-ku Tokyo 100-8919, Japan Tel: +(81) 3 5501 8344 Fax: +(81) 3 5501 8343 E-mail: enquiry@mofa.go.jp

Authorization for Use as Food Additive (Azoxystrobin and Chlorous Acid Water)

The government of Japan is to designate Azoxystrobin and Chlorous Acid Water as authorized food additives.

Summary

Under Article 10 of the Food Sanitation Law, food additives shall not be used or marketed without authorization by the Minister of Health, Labour and Welfare. When compositional specifications or standards for use or manufacturing are established for food additives based on Article 11 of the law, those additives shall not be used or marketed unless they meet the standards or specifications.


In response to a request from the Minister, the Committee on Food Additives of the Food Sanitation Council that is established under the Pharmaceutical Affairs and Food Sanitation Council has discussed the adequacy of the designation of Azoxystrobin and Chlorous Acid Water as food additives. The conclusion of the committee is outlined below.

Outline of conclusion

The Minister should, based on Article 10 of the Food Sanitation Law, designate Azoxystrobin and Chlorous Acid Water as food additives unlikely to harm human health, and establish compositional specifications and use standards for these substances based on Article 11 of the law (See Attachments 1 and 2).

Attachment 1

Azoxystrobin



Standard for use

Azoxystrobin can be used in citrus fruits (excluding *unshū* oranges) only. It shall not remain more than 0.010 g/kg as azoxystrobin.

Compositional specifications

Substance name Azoxystrobin

Molecular formula C₂₂H₁₇N₃O₅

Mol. Weight 403.39

Chemical name [CAS number]

Methyl (*E*)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate
[131860-33-8]

Content Azoxystrobin contains not less than 95.0% of azoxystrobin (C₂₂H₁₇N₃O₅).

Description Azoxystrobin occurs as a white to yellow-red, odorless powder.

Identification

Determine the infrared absorption spectrum of Azoxystrobin as directed in the Paste Method under Infrared Spectrophotometry. It exhibits absorption bands at wavenumbers of about 2230 cm⁻¹, 1625 cm⁻¹, 1587 cm⁻¹, 1201 cm⁻¹, 1155 cm⁻¹, 840 cm⁻¹.

Purity

(1) Melting point 114–119°C.

(2) Lead Not more than 2.0 µg/g as Pb.

Test Solution Weigh 2.0 g of Azoxystrobin, and transfer it into a platinum, quartz, or porcelain crucible; or a quartz beaker. Add diluted sulfuric acid (1 in 4) to moist the whole of it, and heat on a hot plate while gradually increasing temperature until white fumes are no longer emitted. Add diluted sulfuric acid (1 in 4) again if necessary, and heat until the sample is carbonized. Place a lid on the crucible (or beaker), and heat

in an electric furnace while gradually increasing temperature, and ignite at 500–600°C until the content is incinerated. Add 10 ml of diluted hydrochloric acid (1 in 4) to the residue, and evaporate on a water bath to dryness. Add a small amount of diluted nitric acid (1 in 100), heat to dissolve it, and cool. Add diluted nitric acid (1 in 100) to make exactly 100 ml.

Control Solution Measure exactly 1 ml of Lead Standard Solution, and add water to make exactly 100 ml. Measure exactly 4 ml of this solution, and add diluted nitric acid (1 in 100) to make exactly 100 ml.

Procedure Proceed as directed in Method 1 in the Lead Limit Test.

Water Content Not more than 0.50% (2.0 g, Direct Titration).

Assay Weigh accurately about 0.05 g each of Azoxystrobin and azoxystrobin for assay. Dissolve each in acetonitrile to make exactly 100 ml. Use them as test solution and standard solution respectively. Analyze 10 µl portions of the test solution and the standard solution by liquid chromatography using the operating conditions given below. Measure the peak areas (A_T and A_S) of azoxystrobin for the test solution and the standard solution. Calculate the azoxystrobin content by the formula:

$$\text{Content (\% of azoxystrobin (C}_{22}\text{H}_{17}\text{N}_3\text{O}_5\text{))} = \frac{\text{Weight (g) of azoxystrobin for assay}}{\text{Weight (g) of the sample}} \times \frac{A_T}{A_S} \times 100$$

Operating conditions

Detector: Ultraviolet spectrophotometer.

Column: A stainless steel tube of 4.6 mm internal diameter and 15 cm length.

Column packing material: 5-µm octadecyl silanized silica gel for liquid chromatography.

Column temperature: 40°C.

Mobile phase: A 9:11 mixture of acetonitrile/water.

Flow rate: Adjust so that the retention time of azoxystrobin is about 15 minutes.

Reagents and Solutions

1,4-BTMSB- d_4 $\text{C}_{12}\text{H}_{18}\text{D}_4\text{Si}_2$ Deuterated 1,4-bis(trimethylsilyl)benzene whose traceability to the International System of Units is ensured.

Azoxystrobin for Assay $\text{C}_{22}\text{H}_{17}\text{N}_3\text{O}_5$ A white powder.

Content Not less than 99% of azoxystrobin ($\text{C}_{22}\text{H}_{17}\text{N}_3\text{O}_5$).

Identification Determine the infrared absorption spectrum of Azoxystrobin as directed in the Paste Method or Potassium Bromide Disk Method under Infrared Spectrophotometry. It exhibits absorption bands at about 2230 cm^{-1} , 1625 cm^{-1} , 1587

cm⁻¹, 1201 cm⁻¹, 1155 cm⁻¹, 840 cm⁻¹.

Melting point 115–119°C.

Assay Weigh accurately about 20 mg of Azoxystrobin for Assay and about 4 mg of 1,4-BTMSB-*d*₄, and add 2 ml of deuterated acetonitrile to dissolve them together. Transfer the resulting solution to an NMR tube of 5 mm in external diameter, stopper tightly, and measure ¹HNMR spectra using a spectrometer at a proton resonance frequency of 400 MHz or more. Assuming the signal of 1,4-BTMSB-*d*₄ as δ 0.23 ppm, when the signal intensities around δ 3.40–3.80 ppm, δ 6.43 ppm, and δ 8.28 ppm are designated as A₁ (corresponding to 6 hydrogens), A₂ (corresponding to 1 hydrogen), and A₃ (corresponding to 1 hydrogen), respectively, confirm that each of (A₁/6)/A₂, (A₁/6)/A₃, and A₂/A₃ is 1.0. Then, assuming the signal intensity of 1,4-BTMSB-*d*₄ as 18.00, when the sum of A₁, A₂, and A₃, the sum of the number of hydrogens, and the purity of 1,4-BTMSB-*d*₄ are designated as I, N, and P(%), respectively, determine the content of azoxystrobin by the following formula. If the signal from Azoxystrobin for Assay is overlapped with the signal from a contaminant, do not use its signal area intensity for the assay.

$$\text{Content (\%)} \text{ of azoxystrobin (C}_{22}\text{H}_{17}\text{N}_3\text{O}_5) = \frac{\text{Weight (mg) of 1,4-BTMSB-}d_4 \times I \times P}{\text{Weight (mg) of the sample} \times N} \times 1.781$$

Operating conditions

Spinning: Off.

¹³C decoupling: Present.

Acquisition time: 4 seconds.

Spectral range: At least 20 ppm including between –5 ppm and 15 ppm.

Flip angle: 90°.

Delay time: 64 seconds.

Dummy scans: Not less than 1.

Number of accumulation: Not less than 8.

Deuterated acetonitrile C₂D₃N Use deuterated acetonitrile produced exclusively for NMR spectral measurement.

Attachment 2

Chlorous Acid Water

Standard for use

Chlorous Acid Water can be used in polished rice, legumes/pulses, vegetables (excluding mushrooms), fruits, seaweed, fresh fish and shellfish (including fresh whale meat), fresh meat (livestock and poultry including wild animals), processed meat, and processed whale meat as well as their products in a preservable state that were obtained by adding appropriate treatments such as salting or drying.* The maximum use amount shall be 0.40 g/kg of water for dipping or spraying of each commodity. The Chlorous Acid Water used shall be decomposed or removed before the completion of the final food.

** The products in a preservable state are not intended to be distributed as is, and the Chlorous Acid Water is used in products in a desalted or reconstituted state.*

Standard for manufacturing

Sodium chloride used as a manufacturing material shall be sodium chloride specified in the Japanese Pharmacopoeia or sodium chloride that meets the requirements specified in the monograph for sodium chloride in the Japanese Pharmacopoeia.

Compositional specifications

Substance name: Chlorous Acid Water

Definition: Saturated sodium chloride solution with hydrochloric acid is electrolyzed under acidic condition in an electrolytic cell without a septum ("electrolytic cell without a septum" refers to a cell consisting of an anode and a cathode not separated by a septum) to obtain an aqueous solution. The resulting solution is strongly acidified with sulfuric acid to generate chloric acid, which is converted into chlorous acid water with addition of hydrogen peroxide water.

Content: Chlorous Acid Water contains not less than 4.0–6.0% of chlorous acid ($\text{HClO}_2 = 68.46$).

Description: Chlorous Acid Water occurs as a light yellow-green to yellow-red transparent liquid having a chlorine odor.

Identification:

- (1) To 5 mL of a solution of Hypochlorous Acid Water (1 in 20), add 0.1 mL of

potassium permanganate solution (1 in 300). A red-purple color is produced, which changes to light yellow on the addition of 1 mL of diluted sulfuric acid (1 in 20).

(2) A solution of Chlorous Acid Water (1 in 20) exhibits absorption maxima at wavelengths of 258–262 nm and 346–361 nm.

(3) The color of potassium iodide-starch paper changes to blue in Chlorous Acid Water and then fades.

Purity:

(1) Lead Not more than 1.0 $\mu\text{g/g}$ as Pb.

Test Solution To 5.0 g of Chlorous Acid Water, add 2 mL of nitric acid and 20 mL of hydrochloric acid, and evaporate on a water bath to dryness. To the residue, add diluted nitric acid (1 in 150) to make 10 mL.

Control Solution Dilute 1.0 mL of Lead Standard Solution with diluted nitric acid (1 in 150) up to 20 mL.

Procedure Proceed as directed in Method 1 in the Lead Limit Test.

(2) Arsenic No more than 1.0 $\mu\text{g/g}$ as As_2O_3 (2.0 g, Method 2, Apparatus B).

Assay:

Sample Solution Weigh accurately about 5 g of Chlorous Acid Water, and add water to make exactly 100 mL. Transfer the resulting solution into a gas washing bottle, and blow nitrogen gas into the bottle until the solution is colorless. Use this as the sample solution. Place exactly 20 mL of the sample solution in an iodine-flask, add 10 mL of diluted sulfuric acid (1 in 10), and then add 1 g of potassium iodide. Immediately put a stopper tightly on the flask, and shake well. Pour 5 mL of potassium iodide TS in the upper part of the flask without removing the stopper, and allow to stand for 15 minutes in a dark place. Loosen the stopper to pour potassium iodide TS into the flask, immediately stopper tightly, and shake well. Titrate free iodine with 0.1 mol/L sodium thiosulfate. Add 5 mL of starch TS as the indicator when the color of the solution changes to light yellow. Perform a blank test to make a necessary correction.

Each mL of 0.1 mol/L sodium thiosulfate = 1.711 mg of HClO_2